arXiv:1501.05487v3 [math.GR] 24 Jun 2015

A COMBINATORIAL NEGATIVE CURVATURE CONDITION IMPLYING GROMOV HYPERBOLICITY

IOANA-CLAUDIA LAZĂR
'POLITEHNICA' UNIVERSITY OF TIMIŞOARA, DEPT. OF MATHEMATICS,
VICTORIEI SQUARE 2, 300006-TIMIŞOARA, ROMANIA
E-MAIL ADDRESS: IOANA.LAZAR@UPT.RO

Abstract. We explore a local combinatorial condition on a simplicial complex, called 8–location. We show that 8–location and simple connectivity imply Gromov hyperbolicity. We mention some applications of such combinatorial negative curvature condition.

2010 Mathematics Subject Classification: 05C99, 05C75.

Keywords: 8-location, SD' property, Gromov hyperbolicity, universal cover.

1. Introduction

Curvature can be expressed both in metric and combinatorial terms. Metrically, one can refer to nonpositively curved (respectively, negatively curved) metric spaces in the sense of Aleksandrov, i.e. by comparing small triangles in the space with triangles in the Euclidean plane (hyperbolic plane). These are the CAT(0) (respectively, CAT(-1)) spaces. Combinatorially, one looks for local combinatorial conditions implying some global features typical for nonpositively curved metric spaces.

A very important combinatorial condition of this type was formulated by Gromov [Gro87] for cubical complexes, i.e. cellular complexes with cells being cubes. Namely, simply connected cubical complexes with links (that can be thought as small spheres around vertices) being flag (respectively, 5–large, i.e. flag-no-square) simplicial complexes carry a canonical CAT(0) (respectively, CAT(-1)) metric. Another important local combinatorial condition is local k–largeness, introduced by Januszkiewicz-Świątkowski [JŚ06] and Haglund [Hag03]. A flag simplicial complex is *locally k–large* if its links do not contain 'essential' loops of length less than k. In particular, simply connected locally 7–large simplicial complexes, i.e. 7–*systolic* complexes, are Gromov hyperbolic [JŚ06]. The theory of 7–*systolic groups*, that is groups acting geometrically on 7–systolic complexes, allowed to provide important examples of highly dimensional Gromov hyperbolic groups [JŚ03, JŚ06, Osa13a, OŚ15].

However, for groups acting geometrically on CAT(-1) cubical complexes or on 7–systolic complexes, some very restrictive limitations are known. For example, 7–systolic groups are in a sense 'asymptotically hereditarily aspherical', i.e. asymptotically they can not contain essential spheres. This yields in particular that such groups are not fundamental groups of negatively curved manifolds of dimension above two; see e.g. [JŚ07, Osa07, Osa08, OŚ15, GO14, Osa15b]. This rises need for other combinatorial conditions, not imposing restrictions as above. In

[Osa13b, CO15, BCC$^+$13, CCHO14] some conditions of this type are studied – they form a way of unifying CAT(0) cubical and systolic theories. On the other hand, Osajda [Osa15a] introduced a local combinatorial condition of 8–*location*, and used it to provide a new solution to Thurston's problem about hyperbolicity of some 3–manifolds.

In the current paper we undertake a systematic study of a version of 8–location, suggested in [Osa15a, Subsection 5.1]. This version is in a sense more natural than the original one (tailored to Thurston's problem), and none of them is implied by the other. However, in the new 8–location we do allow essential 4–loops. This suggests that it can be used in a much wider context. Roughly (see Section 2 for the precise definition), the new 8–location says that essential loops of length at most 8 admit filling diagrams with at most one internal vertex.

We show that this local combinatorial condition is a negative-curvature-type condition, by proving the following main result of this paper.

Theorem 1.1. *Let X be a simply connected, 8–located simplicial complex. Then the 1–skeleton of X, equipped with the standard path metric, is Gromov hyperbolic.*

The above theorem was announced without a proof in [Osa15a, Subsection 5.1]. In [Osa15a, Subsection 5.2] applications concerning some weakly systolic complexes and groups are mentioned.

Our proof consists of two steps. In Theorem 3.2 we show that an 8–located simplicial complex satisfying a global condition SD' (see Definition 2.2) is Gromov hyperbolic. Then, in Theorem 4.1 we show that the universal cover of an 8–located complex satisfies the property SD'. The main Theorem 4.3 follows immediately from those two results. For proving Theorem 4.1 we use a method of constructing the universal cover introduced in [Osa13b], and then developed in [BCC$^+$13, CCHO14].

Acknowledgements. I am indebted to Damian Osajda for introducing me to the subject, posing the problem, helpful explanations, and patience. I thank Louis Funar for introducing me to systolic geometry a few years ago. I would also like to thank the Mathematical Institute in Wrocław, Poland for hospitality during the winter of 2014. The visit was partially supported by (Polish) Narodowe Centrum Nauki, decision no DEC-2012/06/A/ST1/00259.

2. Preliminaries

Let X be a simplicial complex. We denote by $X^{(k)}$ the k-skeleton of $X, 0 \leq k < \dim X$. A subcomplex L in X is called *full* (in X) if any simplex of X spanned by a set of vertices in L, is a simplex of L. For a set $A = \{v_1, ..., v_k\}$ of vertices of X, by $\langle A \rangle$ or by $\langle v_1, ..., v_k \rangle$ we denote the *span* of A, i.e. the smallest full subcomplex of X that contains A. We write $v \sim v'$ if $\langle v, v' \rangle \in X$ (it can happen that $v = v'$). We write $v \nsim v'$ if $\langle v, v' \rangle \notin X$. We call X *flag* if any finite set of vertices, which are pairwise connected by edges of X, spans a simplex of X.

A *cycle* (*loop*) γ in X is a subcomplex of X isomorphic to a triangulation of S^1. A *k-wheel* in X $(v_0; v_1, ..., v_k)$ (where $v_i, i \in \{0, ..., k\}$ are vertices of X) is a subcomplex of X such that $(v_1, ..., v_k)$ is a full cycle and $v_0 \sim v_1, ..., v_k$. The *length* of γ (denoted by $|\gamma|$) is the number of edges in γ.

Definition 2.1. A simplicial complex is *m-located* if it is flag and every full homotopically trivial loop of length at most m is contained in a 1-ball.

The *link* of X at σ, denoted X_σ, is the subcomplex of X consisting of all simplices of X which are disjoint from σ and which, together with σ, span a simplex of X. A *full cycle* in X is a cycle that is full as subcomplex of X. We call a flag simplicial complex *k-large* if there are no full j-cycles in X, for $j < k$. We say X is *locally k-large* if all its links are k-large.

We define the *metric* on the 0-skeleton of X as the number of edges in the shortest 1−skeleton path joining two given vertices and we denote it by d. A *ball (sphere)* $B_i(v, X)$ ($S_i(v, X)$) of radius i around some vertex v is a full subcomplex of X spanned by vertices at distance at most i (at distance i) from v.

We introduce further a global combinatorial condition on a flag simplicial complex.

Definition 2.2. Let X be a flag simplicial complex. For a vertex O of X and a natural number n, we say that X satisfies *the property $SD'_n(O)$* if for every $i \in \{1, ..., n\}$ we have:

1. (T) (triangle condition): for every edge $e \in S_{i+1}(0)$, the intersection $X_e \cap B_i(O)$ is non-empty;
2. (V) (vertex condition): for every vertex $v \in S_{i+1}(0)$, and for every two vertices $u, w \in X_v \cap B_i(O)$, there exists a vertex $t \in X_v \cap B_i(O)$ such that $t \sim u, w$.

We say X satisfies *the property $SD'(O)$* if $SD'_n(O)$ holds for each natural number n. We say X satisfies *the property SD'* if $SD'_n(O)$ holds for each natural number n and for each vertex O of X.

The following result is given in [Osa15a].

Proposition 2.1. *A simplicial complex which satisfies the property $SD'(O)$ for some vertex O, is simply connected.*

By a *covering* we mean a *simplicial covering*, i.e. a simplicial map restricting to isomorphisms from 1-balls onto their images.

3. HYPERBOLICITY

In this section we show that the 8-location on a simplicial complex enjoying the SD' property, implies Gromov hyperbolicity.

Lemma 3.1. *Let X be an 8-located simplicial complex which satisfies the $SD'_n(O)$ property for some vertex O. Let $v \in S_{n+1}(O)$ and let $y, z \in B_n(O)$ be such that $v \sim y, z$ and $d(y, z) = 2$. Let $w \in B_n(O)$ be a vertex such that $w \sim y, v, z$, given by the vertex condition (V). Consider the vertices $u_1, u_2 \in B_{n-1}(O)$ such that $\langle y, u_1, w\rangle, \langle w, u_2, z\rangle \in X$, given by the triangle condition (T) and such that $u_1 \nsim z$ and $u_2 \nsim y$. Then $u_1 \sim u_2$ (possibly with $u_1 = u_2$).*

Proof. The proof is by contradiction. Assume that $d(u_1, u_2) = 2$. Let $u' \in B_{n-1}(O)$ be a vertex such that $u' \sim u_1, w, u_2$ given by the vertex condition (V). Let $t_1, t_2 \in B_{n-2}(O)$ be vertices such that $\langle u_1, t_1, u'\rangle, \langle u', t_2, u_2\rangle \in X$ given by the triangle condition (T). Let $t' \in B_{n-2}(O)$ be a vertex such that $t' \sim t_1, u', t_2$ (possibly with $t' = t_2$) given by the vertex condition (V). Note that if $u_1 \nsim t_2$ and $u_2 \nsim t_1$, the full homotopically trivial loop $(v, z, u_2, t_2, t', t_1, u_1, y)$ has length at most 8. If $u_1 \sim t'$, consider the full homotopically trivial loop $(v, z, u_2, t_2, t', u_1, y)$ of length at most 7. If $u_1 \sim t_2$ or $t_1 = t_2$, consider the full homotopically trivial loop (v, z, u_2, t_2, u_1, y) of length 6. In all three cases it follows, by 8-location, that $d(v, t_2) = 2$. But since $v \in S_{n+1}$ whereas $t_2 \in S_{n-2}$, we have $d(v, t_2) = 3$. Hence, because we have reached a contradiction, it follows that $u_1 \sim u_2$ (possibly with $u_1 = u_2$).



$\square$

Theorem 3.2. *Let X be an 8-located simplicial complex which satisfies the SD' property. Then the $0-$skeleton of X with a path metric induced from $X^{(1)}$, is $\delta-$hyperbolic, for a universal constant δ.*

Proof. The proof is similar to the one of the analogous Theorem 3.3 given in [Osa15a]. According to [Pap95], we can prove hyperbolicity of the 0-skeleton by showing that intervals are uniformly thin.

Let O, O' be two vertices. Denote by I the set of vertices lying on geodesics between O and O' and let $n = d(O, O')$. Let I_k denote the intersection $S_k(O) \cap S_{n-k}(O') = S_k(O) \cap I$. We prove by contradiction that for every $k \leq n$ and for every two vertices $v, w \in I_k, d(v, w) \leq 2$. This also shows that the hyperbolicity constant is universal.

We build a full path of diameter 3 as in [Osa15a]. Suppose there are vertices $v, w \in I_k$ such that $d(v, w) > 2$. Let k be the maximal natural number for which this happens. Then there exist vertices v', w' in I_{k+1} such that $v' \sim v$, $w' \sim w$, and $d(v', w') \leq 2$.

By the vertex condition (V), there is a vertex z in I_{k+1} such that $z \sim v', w'$, possibly with $z = w'$. By the triangle condition (T) there are vertices $v'', w'' \in I_k$ such that $\langle v', v'', z\rangle, \langle z, w'', w'\rangle \in X$ (with $v'' = w''$ if $z = w'$). By the vertex condition (V) there are vertices s, t and u in I_k such that $s \sim v, v', v''; t \sim v'', z, w''; u \sim w, w', w''$ (possibly with $s = v'', t = w''$, and $u = w$). Among the vertices t, w'', u, w we choose the first one (in the given order), that is at distance 3 from v. Denote this vertex by v'''. In this way we obtain a full path (v_1, v_2, v_3, v_4) in I_k of diameter 3, with $v_1 = v$ and $v_4 = v'''$. We will show that such a path can not exist reaching hereby a contradiction and proving the theorem.

By the triangle condition (T) there exist vertices w_i in I_{k-1} such that $\langle v_i, w_i, v_{i+1}\rangle \in X, 1 \leq i \leq 3$. We discuss further all possible cases (and the corresponding subcases) of mutual relations between vertices $w_i, 1 \leq i \leq 3$ (up to renaming vertices). Case I is when $w_1 = w_2$ and $w_2 \neq w_3$. Case II is when $w_1 \neq w_2 \neq v_3$.

By the triangle condition (T) there are vertices p_i in I_{k+1} such that $\langle v_i, p_i, v_{i+1}\rangle \in X, 1 \leq i \leq 3$ (possibly with $p_2 = p_3$). By the vertex condition (V) there exist vertices p', p'' in I_{k+1} such that $p' \sim p_1, v_2, p_2$ and $p'' \sim p_2, v_3, p_3$ (possibly with $p' = p_2$ and $p'' = p_3$).

Note that if $p_1 = p_3$ or $w_1 = w_3$, then $d(v_1, v_4) = 2$ which yields a contradiction because $d(v_1, v_4) = 3$.

3.1. **Case I.** We start treating the case when $w_1 = w_2$ and $w_2 \neq w_3$ which has 2 subcases. Case I.1 is when $w_1 = w_2, w_2 \sim w_3$. Case I.2 is when $w_1 = w_2, d(w_2, w_3) = 2$.

According to Lemma 3.1, since $d(v_1, v_3) = 2$, we get $d(p_1, p_2) \leq 1$.

3.1.1. *Case I.1.* We treat the case when $w_1 = w_2, w_2 \sim w_3$.

Claim 3.3. *Among the vertices p_1, p_2, p'', p_3 in this order we choose the last vertex p^* such that $v_1 \sim p^*$. Among the remaining vertices in the same order, namely among the sequence of vertices starting with p^* and ending with p_3, we choose the first vertex p^{**} such that $p^{**} \sim v_4$. Let γ be the homotopically trivial loop containing at least the vertices $p^{**}, v_4, w_3, w_2, v_1, p^*$. If $p^* \sim p^{**}$ then $\gamma = (p^{**}, v_4, w_3, w_2, v_1, p^*)$. If $p^* \neq p_2$, $p^{**} \neq p_2$ and $p^* \nsim p^{**}$ then $p_2 \in \gamma$. If $p^* \neq p''$, $p^{**} \neq p''$ and $p^* \nsim p^{**}$ then $p'' \in \gamma$. Then the loop γ is full.*

Proof. Note that $p^* \neq p^{**}$ since otherwise we get contradiction with $d(v_1, v_4) = 3$. We also get contradiction with $d(v_1, v_4) = 3$ if $v_1 \sim w_3$ or $v_4 \sim w_2$. Hence, due to the choice of the vertices p^* and p^{**}, the cycle γ is full.

□

In order to apply 8-location, we consider a homotopically trivial loop γ containing a subset of the set of vertices of the cycle $(p_3, v_4, w_3, w_2, v_1, p_1, p_2, p'')$, or all vertices of this cycle. Namely, γ contains the vertices v_4, w_3, w_2, v_1 and at least two of the vertices p_1, p_2, p_3, p'' chosen as in Claim 3.3. Note that γ is full. Thus, since its length is at most 8, γ is contained in the link of a vertex. So we get $d(v_1, v_4) = 2$ which yields contradiction with $d(v_1, v_4) = 3$.



Case I.1

3.1.2. *Case I.2.* We treat further the case when $w_1 = w_2, d(w_2, w_3) = 2$. By the vertex condition (V) there exists a vertex $w' \in I_{k+1}$ such that $w' \sim w_2, v_3, w_3$ (possibly with $w' = w_3$).

We consider the homotopically trivial loop $(p_3, v_4, w_3, w', w_2, v_1, p_1, p_2, p'')$ of length at most 9. Note that there are no edges between the vertices w_2, w_3, w' and p_1, p_2, p_3, p''. But there can be diagonals between the vertices v_1, v_2, v_3, v_4 and p_1, p_2, p_3, p''. Also there can be diagonals joining the vertices w_2, w_3, w' and v_1, v_2, v_3, v_4. Case I.2.a is when such diagonals exist, or some of the vertices p_1, p_2, p'', p_3 coincide, or nonconsecutive vertices in this sequence are adjacent. Case I.2.b is when none of these situations occur.



Case I.2.b

Note that if $v_1 \sim w'$ and $v_4 \sim w'$, or if $v_1 \sim w_3$, or if $v_4 \sim w_2$, we have contradiction with $d(v_1, v_4) = 3$. If $v_1 \sim w'$ but $v_4 \nsim w'$, or if $v_4 \sim w'$ but $v_1 \nsim w'$, we have reached case I.1 treated above.

Note that if $v_2 \sim w_3$ Lemma 3.1 implies $d(p_2, p_3) \leq 1$. We choose a loop β as in Claim 3.3 containing at most all vertices of $\gamma = (p_3, v_4, w_3, w', w_2, v_1, p_1, p_2)$. Namely, β contains the vertices v_1, v_4, at least two of the vertices p_1, p_2, p_3, and at least two of the vertices w_2, w_3, w'. Because β is full and has length at most 8, by 8-location we get contradiction with $d(v_1, v_4) = 3$.

In case I.2.a we consider a homotopically trivial loop γ containing a subset of the set of vertices of the cycle $(p_3, v_4, w_3, w', w_2, v_1, p_1, p_2, p'')$ chosen as in Claim 3.3, or all of its vertices. Namely, γ contains the vertices v_4, v_1, at least two of the vertices p_1, p_2, p_3, p'', and at least two of the vertices w_2, w_3, w'. Note that γ is full. In order to apply 8-location, the length of γ should be at most 8. If so, since γ is contained, by 8-location, in the link of a vertex, we get contradiction with $d(v_1, v_4) = 3$.

For the rest of case I.2.a, we consider the situation when there is no full homotopically trivial loop γ as in the paragraph above of length at most 8. This happens if $d(p_1, p_2) + d(p_2, p_3) = 3$ (if $d(p_1, p_2) + d(p_2, p_3) \leq 2$, the length of γ is at most 8) and if there are no diagonals between the vertices v_1, v_4 and the vertices p_1, p_2, p_3, p'' (if such diagonals exist, we can find a full loop γ as in the paragraph above of length at most 8). But because we are still in case I.2.a, either there exists at least one diagonal between the vertices v_2, v_3 and the vertices p_1, p_2, p_3, p'', or nonconsecutive vertices in the sequence p_1, p_2, p'', p_3 are adjacent, or both.

- Consider first the case $d(p_1, p_2) + d(p_2, p_3) = 3$, $p_1 \nsim v_3, p_1 \nsim p''$. By the triangle condition (T) there are vertices q_1, q_2 in I_{k+2} such that $\langle p_1, q_1, p_2\rangle, \langle p_2, q_2, p''\rangle \in X$. By the vertex condition (V) there is a vertex q' in I_{k+2} such that $q' \sim q_1, p_2, q_2$ (possibly with $q' = q_2$). Note that if $p'' \sim v_2$, Lemma 3.1 implies that $d(q_1, q_2) \leq 1$. Consider a homotopically trivial loop γ containing a subset of the set of vertices of the cycle $(q_2, p'', v_3, w_2, v_1, p_1, q_1, q')$ chosen as in Claim 3.3, or all of its vertices. Namely, γ contains the vertices p'', v_3, w_2, v_1, p_1, and at least one of the vertices q_1, q_2, q' chosen as in Claim 3.3 (note that here the vertices p^* and p^{**} as denoted in Claim 3.3 may coincide). Because γ is full and has length at most 8, by 8-location we get contradiction with $d(q_1, w_2) = 3$.

- Consider further the case $d(p_1, p_2) + d(p_2, p_3) = 3$, $p_1 \nsim v_3, p_1 \sim p''$. By the triangle condition (T) there are vertices q_1, q_2 in I_{k+2} such that $\langle p_1, q_1, p''\rangle, \langle p'', q_2, p_3\rangle \in X$. By the vertex condition (V) there is a vertex q' in I_{k+2} such that $q' \sim q_1, p'', q_2$ (possibly with $q' = q_2$). Consider a homotopically trivial loop γ containing a subset of the set of vertices of the cycle $(q_2, p_3, v_3, w_2, v_1, p_1, q_1, q')$ chosen as in Claim 3.3, or all of its vertices. Namely, γ contains the vertices p_3, v_3, w_2, v_1, p_1, and at least



Case I.2.a: $d(p_1, p_2) + d(p_2, p_3) = 3, p_1 \nsim v_3, p_1 \nsim p''$

one of the vertices q_1, q_2, q' chosen as in Claim 3.3. Because γ is full and has length at most 8, by 8-location we get contradiction with $d(q_1, w_2) = 3$.



Case I.2.a: $d(p_1, p_2) + d(p_2, p_3) = 3, p_1 \nsim v_3, p_1 \sim p''$

- Consider further the situation when $d(p_1, p_2) + d(p_2, p_3) = 3$ and $p_1 \sim v_3$. By the triangle condition (T) there are vertices $q_1, q_2, q_3 \in I_{k+2}$ such that $\langle p_1, q_1, p_2\rangle$, $\langle p_2, q_2, p''\rangle, \langle p'', q_3, p_3\rangle \in X$. Note that according to Lemma 3.1, $d(q_2, q_3) \leq 1$. By the vertex condition (V) there exists a vertex $q' \in I_{k+2}$ such that $q' \sim q_1, p_2, q_2$ (possibly with $q' = q_2$). In order to apply 8-location, we consider a homotopically trivial loop γ containing a subset of the set of vertices of the cycle $(q_3, p_3, v_3, p_1, q_1, q', q_2)$ chosen as in Claim 3.3, or all vertices of this cycle. Namely, γ contains the vertices p_1, v_3, p_3, and at least one of the vertices q_1, q_2, q_3, q'. Since γ is full, by 8-location, there exists a vertex p_{02} adjacent to all vertices of γ. Note that $p_{02} \in I_{k+1}$. We choose a homotopically trivial loop as in Claim 3.3 containing the vertices v_1, v_4, at least two of the vertices p_1, p_{02}, p_3, and at least two of the vertices w_2, w_3, w'. By 8-location, we obtain contradiction with $d(v_1, v_4) = 3$.



Case I.2.a: $d(p_1, p_2) + d(p_2, p_3) = 3, p_1 \sim v_3$

Note that case I.2.b is similar to case $d(p_1,p_2)+d(p_2,p_3)=3$, $p_1 \nsim v_3, p_1 \nsim p''$ treated above.

3.2. **Case II.** We treat next the case when $w_1 \neq w_2 \neq w_3$ which has 3 subcases. Case II.1 is when $w_1 \sim w_2 \sim w_3$. Case II.2 is when $w_1 \sim w_2$ and $d(w_2,w_3)=2$. Case II.3 is when $d(w_1,w_2)=d(w_2,w_3)=2$.

Note that if $w_1 \sim v_3$, Lemma 3.1 implies that $d(p_1,p_2)\leq 1$. Also note that if $w_3 \sim v_2$, Lemma 3.1 implies that $d(p_2,p_3)\leq 1$. So in case $w_1 \sim v_3$ and $w_3 \sim v_2$, we choose a loop γ containing at most all vertices of the homotopically trivial $(p_3, v_4, w_3, w_2, w_1, v_1, p_1, p_2)$. Namely, γ contains the vertices v_1, v_4, at least two of the vertices p_1, p_2, p_3, and at least two of the vertices w_1, w_2, w_3. Because γ is full and has length at most 8, we get, by $8-$location, contradiction with $d(v_1,v_4)=3$.

We consider the situation when $w_1 \sim w_3$. We consider a homotopically trivial loop γ containing at most all vertices of the cycle $(p_2, v_3, w_3, w_1, v_1, p_1, p')$ chosen as in Claim 3.3. Namely, γ contains the vertices v_3, w_3, w_1, v_1, and at least one of the vertices p_1, p_2, p'. Because γ is full and has length at most 7, it is contained, by 8-location, in the link of a vertex v_{02}. Note that $v_{02} \in I_k$. Hence, because the path (v_1, v_2, v_3, v_4) is full, and $v_{02} \sim v_1, v_3$, the path (v_1, v_{02}, v_3, v_4) is also full. Moreover, since $\langle v_1, w_1, v_{02}\rangle$, $\langle v_{02}, w_3, v_3\rangle$, $\langle v_3, w_3, v_4\rangle \in X$ and $w_1 \sim w_3$, we have reached case I.1 treated above.



Case II: $w_1 \sim w_3$

3.2.1. *Case II.*1. We treat the case when $w_1 \sim w_2 \sim w_3$.

We consider the homotopically trivial loop $(p_3, v_4, w_3, w_2, w_1, v_1, p_1, p', p_2, p'')$ of length at most 10. Note that there are no diagonals between the vertices w_1, w_2, w_3 and p_1, p_2, p_3, p', p''. But there can be diagonals joining the vertices w_1, w_2, w_3 to the vertices v_1, v_2, v_3, v_4. Also there can be diagonals between the vertices v_1, v_2, v_3, v_4 and p_1, p_2, p_3, p', p''. Case II.1.a is when such diagonals exist, or some of the vertices p_1, p', p_2, p'', p_3 coincide, or nonconsecutive vertices in this sequence are adjacent. Case II.1.b is when none of these situations occur.

Note that if $v_1 \sim w_2$ and $v_4 \sim w_2$, or if $v_1 \sim w_3$, or if $v_4 \sim w_1$, we have contradiction with $d(v_1,v_4)=3$. If $v_1 \sim w_2$ but $v_4 \nsim w_2$, or if $v_4 \sim w_2$ but $v_1 \nsim w_2$, we have reached case I.1.

In case II.1.a, we consider a homotopically trivial loop γ containing a subset of the set of vertices of the cycle $(p_3, v_4, w_3, w_2, w_1, v_1, p_1, p', p_2, p'')$ chosen as in Claim



Case II.1.b

3.3, or all of its vertices. Namely, γ contains the vertices v_4, v_1, at least two of the vertices p_1, p_2, p_3, p', p'' chosen as in Claim 3.3, and at least two of the vertices w_1, w_2, w_3 chosen as in Claim 3.3. Note that γ is full. In order to apply 8-location, the length of γ should be at most 8. If so, since γ is contained, by 8-location, in the link of a vertex, we get contradiction with $d(v_1, v_4) = 3$.

For the rest of case II.1.a, we consider the situation when there is no full homotopically trivial loop γ as in the paragraph above of length at most 8. This happens if $d(p_1, p_2) + d(p_2, p_3) \geq 3$ (if $d(p_1, p_2) + d(p_2, p_3) \leq 2$, the length of γ is at most 8) and if there are no diagonals between the vertices v_1, v_4 and the vertices p_1, p_2, p_3, p', p'' such that we can find a full loop γ as in the paragraph above of length at most 8. Because we are still in case II.1.a, either one, or some of the following situations occur. Namely, either there exists at least one diagonal between the vertices v_1, v_4 and the vertices p_1, p_2, p_3, p', p'' such that we can find a full loop γ as in the paragraph above of length greater than 8, or else nonconsecutive vertices in the sequence p_1, p', p_2, p'', p_3 are adjacent, or else there exists at least one diagonal between the vertices v_2, v_3 and the vertices p_1, p_2, p_3, p', p''.

- We treat first the case $d(p_1, p_2) = d(p_2, p_3) = 2$, $p_1 \nsim v_3$. By the triangle condition (T) there are vertices q_1, q_2 in I_{k+2} such that $\langle p_1, q_1, p' \rangle, \langle p', q_2, p_2 \rangle \in X$. Lemma 3.1 implies that $d(q_1, q_2) \leq 1$. We consider a homotopically trivial loop γ containing at most all vertices of the cycle $(q_2, p_2, v_3, w_2, w_1, v_1, p_1, q_1)$ chosen as in Claim 3.3. Namely, γ contains the vertices p_2, v_3, v_1, p_1, at least one of the vertices q_1, q_2 chosen as in Claim 3.3, and at least one of the vertices w_1, w_2 chosen as in Claim 3.3. Because γ is full and has length at most 8, by 8-location we get contradiction with $d(q_1, w_1) = 3$.



Case II.1.a: $d(p_1, p_2) = d(p_2, p_3) = 2, p_1 \nsim v_3$

Note that if $p_1 \sim p_2$, the loop γ chosen in the previous paragraph is no longer full. This situation will be treated separately in the next two paragraphs. Note that the case when there exists a diagonal between the vertices v_1, v_4 and the vertices

p_1, p_2, p_3, p', p'' such that we can find a full loop γ of length 9 containing the vertices v_4, w_3, w_2, w_1, v_1 and 4 of the vertices p_1, p_2, p_3, p', p'', can be treated similarly.

- We treat further the case $d(p_1, p_2) = 1, d(p_2, p_3) = 2$, $p_3 \nsim v_2$. By the triangle condition (T) there are vertices $q_2, q_3 \in I_{k+2}$ such that $\langle p_2, q_2, p'' \rangle, \langle p'', q_3, p_3 \rangle \in X$. By Lemma 3.1 we get $d(q_2, q_3) \leq 1$. We consider a homotopically trivial loop γ containing a subset of the set of vertices of the cycle $(q_3, p_3, v_4, w_3, w_2, v_2, p_2, q_2)$ chosen as in Claim 3.3, or all of its vertices. Namely, γ contains the vertices p_3, v_4, v_2, p_2, at least one of the vertices q_2, q_3, and at least one of the vertices w_2, w_3. Because γ is full and has length at most 8, by 8-location we get contradiction with $d(q_2, w_2) = 3$.



Case II.1.a: $d(p_1, p_2) = 1, d(p_2, p_3) = 2, p_3 \nsim v_2$

- We treat further the case $d(p_1, p_2) = 1, d(p_2, p_3) = 2$, $p_3 \sim v_2$. By the triangle condition (T) there are vertices $q_1, q_2, q_3 \in I_{k+2}$ such that $\langle p_1, q_1, p_2 \rangle, \langle p_2, q_2, p'' \rangle$, $\langle p'', q_3, p_3 \rangle \in X$. Note that according to Lemma 3.1, $d(q_2, q_3) \leq 1$. By the vertex condition (V) there exists a vertex $q' \in I_{k+2}$ such that $q' \sim q_1, p_2, q_2$ (possibly with $q' = q_2$). Note that if $v_2 \sim p''$, Lemma 3.1 implies that $d(q_1, q_2) \leq 1$. We consider a homotopically trivial loop γ containing a subset of the set of vertices of the cycle $(q_3, p_3, v_2, p_1, q_1, q', q_2)$ chosen as in Claim 3.3, or all of its vertices. Namely, γ contains the vertices p_3, v_2, p_1, and at least one of the vertices q_1, q_2, q_3, q' chosen as in Claim 3.3. Because γ is full and has length at most 7, by 8-location there is a vertex p_{02} adjacent to all vertices of γ. So $p_{02} \in I_{k+1}$. We choose a homotopically trivial loop as in Claim 3.3 containing the vertices v_1, v_4, at least two of the vertices p_1, p_{02}, p_3, and at least two of the vertices w_1, w_2, w_3. By 8-location, we obtain contradiction with $d(v_1, v_4) = 3$.



Case II.1.a: $d(p_1, p_2) = 1, d(p_2, p_3) = 2, p_3 \sim v_2$

- Consider further the case $d(p_1, p_2) = d(p_2, p_3) = 2, p_1 \sim v_3$. By the triangle condition (T) there are vertices $q_1, q_2, q_3 \in I_{k+2}$ such that $\langle p_1, q_1, p' \rangle, \langle p', q_2, p_2 \rangle$,

$\langle p_2, q_3, p''\rangle \in X$. According to Lemma 3.1, $d(q_1, q_2) \leq 1$. Note that if $v_2 \sim p''$ or/and $v_3 \sim p'$, Lemma 3.1 implies that $d(q_2, q_3) \leq 1$. By the vertex condition (V) there is a vertex $q'' \in I_{k+2}$ such that $q'' \sim q_2, p_2, q_3$ (possibly with $q'' = q_3$). We consider a homotopically trivial loop γ containing a subset of the set of vertices of the cycle $(q_3, p'', v_3, p_1, q_1, q_2, q'', q_3)$ chosen as in Claim 3.3, or all of its vertices. Namely, γ contains the vertices p'', v_3, p_1, and at least one of the vertices q_1, q_2, q_3, q''. Because γ is full and has length at most 7, by 8-location there is a vertex p_{02} adjacent to all vertices of γ. So $p_{02} \in I_{k+1}$. By the triangle condition (T) there are vertices $r_1, r_2, r_3 \in I_{k+2}$ such that $\langle p_1, r_1, p_{02}\rangle, \langle p_{02}, r_2, p''\rangle, \langle p'', r_3, p_3\rangle \in X$. Note that since $p_{02} \sim v_3$, Lemma 3.1 implies that $d(r_2, r_3) \leq 1$. By the vertex condition (V) there exists a vertex $r' \in I_{k+2}$ such that $r' \sim r_1, p_{02}, r_2$ (possibly with $r' = r_2$). Note that if $v_2 \sim p''$ Lemma 3.1 implies that $d(r_1, r_2) \leq 1$. We consider a homotopically trivial loop β containing at most all vertices of the cycle $(r_3, p_3, v_3, p_1, r_1, r', r_2)$ chosen as in Claim 3.3. Namely, β contains the vertices p_3, v_3, p_1, and at least one of the vertices r_1, r_2, r_3, r' chosen as in Claim 3.3. Because β is full and has length at most 7, by 8-location there exists a vertex p_{002} adjacent to all vertices of β. So $p_{002} \in I_{k+1}$. Let α be a cycle containing at most all vertices of the cycle $(p_3, v_4, w_3, w_2, w_1, v_1, p_1, p_{002})$ chosen as in Claim 3.3. Namely, α contains the vertices v_1, v_4, at least two of the vertices w_1, w_2, w_3, and at least two of the vertices p_1, p_{002}, p_3. By 8-location, we get contradiction with $d(v_1, v_4) = 3$.



Case II.1.a: $d(p_1, p_2) = d(p_2, p_3) = 2, p_1 \sim v_3$



Case II.1.a: $d(p_1, p_2) = d(p_2, p_3) = 2, p_1 \sim v_3$

Case II.1.b is similar to case $d(p_1, p_2) = d(p_2, p_3) = 2$, $p_1 \nsim v_3$ treated above.

3.2.2. *Case II.2.* We treat further the case when $w_1 \sim w_2$ and $d(w_2, w_3) = 2$. By the vertex condition (V) there is a vertex w' in I_{k-1} such that $w' \sim w_2, v_3, w_3$ (possibly with $w' = w_3$).

Note that if $v_1 \sim w_2$ and $v_4 \sim w_2$, or if $v_1 \sim w_3$, or if $v_4 \sim w_1$, we have contradiction with $d(v_1, v_4) = 3$. If $v_3 \sim w_1$, or $v_2 \sim w_3$ we have reached case I treated above.

By the triangle condition (T) there are vertices $u_1, u_2, u_3 \in I_{k-2}$ such that $\langle w_1, u_1, w_2\rangle$, $\langle w_2, u_2, w'\rangle$, $\langle w', u_3, w_3\rangle \in X$. Lemma 3.1 implies that $d(u_2, u_3) \leq 1$. By the vertex condition (V) there is a vertex $u' \in I_{k-2}$ such that $u' \sim u_1, w_2, u_2$ (possibly with $u' = u_2$).

We consider a homotopically trivial loop γ containing at most all vertices of the cycle $(v_3, w_3, u_3, u_2, u', u_1, w_1, v_2)$ chosen as in Claim 3.3. Namely, γ contains the vertices w_1, v_2, v_3, w_3, and at least one of the vertices u_1, u_2, u_3, u' chosen as in Claim 3.3. Because γ is full and has length at most 8, 8-location implies that there is a vertex w_{02} adjacent to all vertices of γ. So $w_{02} \in I_{k-1}$. Moreover, since $\langle v_1, w_1, v_2\rangle, \langle v_2, w_{02}, v_3\rangle, \langle v_3, w_3, v_4\rangle \in X$ and $w_1 \sim w_{02} \sim w_3$, we have reached case II.1 treated above.



Case II.2.

3.2.3. *Case II.3.* We treat further the case when $d(w_1, w_2) = d(w_2, w_3) = 2$. By the vertex condition (V) there are vertices w', w'' in I_{k-1} such that $w' \sim w_1, v_2, w_2$, and $w'' \sim w_2, v_3, w_3$ (possibly with $w' = w_2$, and $w'' = w_3$).

Note that if $v_1 \sim w'$ and $v_4 \sim w'$, or if $v_1 \sim w_3$, or if $v_4 \sim w_1$, we have contradiction with $d(v_1, v_4) = 3$. Note that if $v_3 \sim w_1$, or if $v_2 \sim w_3$ we have reached case I treated above.

By the triangle condition (T) there are vertices $u_1, u_2, u_3, u_4 \in I_{k-2}$ such that $\langle w_1, u_1, w'\rangle, \langle w', u_2, w_2\rangle$, $\langle w_2, u_3, w''\rangle, \langle w'', u_4, w_3\rangle \in X$. Lemma 3.1 implies that $d(u_1, u_2) \leq 1$ and $d(u_3, u_4) \leq 1$. By the vertex condition (V) there is a vertex $u' \in I_{k-2}$ such that $u' \sim u_2, w_2, u_3$ (possibly with $u' = u_3$).

Consider first the case $d(u_2, u_3) \leq 1$. We consider a homotopically trivial loop γ containing a subset of the set of vertices of the cycle $(v_3, w_3, u_4, u_3, u_2, u_1, w_1, v_2)$ chosen as in Claim 3.3, or all of its vertices. Namely, γ contains the vertices w_1, v_2, v_3, w_3, and at least one of the vertices u_1, u_2, u_3, u_4. Because γ is full and has length at most 8, 8-location implies that there is a vertex w_{02} adjacent to all vertices of γ. So $w_{02} \in I_{k-1}$. Moreover, since $\langle v_1, w_1, v_2\rangle, \langle v_2, w_{02}, v_3\rangle, \langle v_3, w_3, v_4\rangle \in X$ and $w_1 \sim w_{02} \sim w_3$, we have reached case II.1 treated above.

Consider further the situation when $d(u_2, u_3) = 2$. By the triangle condition (T) there are vertices $t_1, t_2 \in I_{k-3}$ such that $\langle u_2, t_1, u'\rangle, \langle u', t_2, u_3\rangle \in X$. Lemma 3.1 implies that $d(t_1, t_2) \leq 1$. We consider a homotopically trivial loop γ containing



Case II.3.: $d(u_2, u_3) = 1$

at most all vertices of the cycle $(v_3, w'', u_3, t_2, t_1, u_2, w', v_2)$. Namely, γ contains the vertices w'', u_3, u_2, w', at least one of the vertices t_1, t_2 chosen as in Claim 3.3, and at least one of the vertices v_2, v_3. Because γ is full and has length at most 8, 8-location implies contradiction with $d(v_2, t_1) = 3$.



Case II.3.: $d(u_2, u_3) = 2$

□

4. Local-to-global

In this section we consider an 8-located simplicial complex and we construct its universal cover such that it has the SD' property.

Theorem 4.1. *Let X be an 8-located simplicial complex. Then its universal cover $\widetilde{X}$ is a simplicial complex which satisfies the SD' property.*

Proof. The proof is similar to the one of the analogous Theorem 3.4 given in [Osa15a].

We construct the universal cover $\widetilde{X}$ of X as an increasing union $\cup_{i=1}^{\infty} \widetilde{B}_i$ of combinatorial balls. The covering map is then the union $\cup_{i=1}^{\infty} f_i : \cup_{i=1}^{\infty} \widetilde{B}_i \to X$, where $f_i : \widetilde{B}_i \to X$ is locally injective and $f_i|_{\widetilde{B}_j} = f_j$, for $j \leq i$.

The proof is by induction. We choose a vertex O of X and we define $\widetilde{B}_0 = \{0\}, \widetilde{B}_1 = B_1(O, X)$, and $f_1 = Id_{B_1(O)}$. We assume that we have constructed the balls $\widetilde{B}_1, \widetilde{B}_2, ..., \widetilde{B}_i$ and the corresponding maps $f_1, f_2, ..., f_i$ to X such that the following conditions hold:

1. (P_i): $\widetilde{B}_j = B_j(O, \widetilde{B}_i), j \in \{1, ..., i\}$;
2. (Q_i): $\widetilde{B}_i$ satisfies the property $SD'_{i-1}(O)$;
3. (R_i): $f_i|_{B_1(\widetilde{w}, \widetilde{B}_i)} : B_1(\widetilde{w}, \widetilde{B}_i) \to B_1(f_i(\widetilde{w}), X)$ is an isomorphism onto the span of the image for $\widetilde{w} \in \widetilde{B}_i$, and it is an isomorphism for $\widetilde{w} \in \widetilde{B}_{i-1}$.

Note that (P_1), (Q_1) and (R_1) hold, i.e. that the above conditions are satisfied for $\widetilde{B}_1$ and f_1. We construct further $\widetilde{B}_{i+1}$ and the map $f_{i+1} : \widetilde{B}_{i+1} \to X$. For a simplex $\widetilde{\sigma}$ of $\widetilde{B}_i$, we denote by σ its image $f_i(\widetilde{\sigma})$ in X. Let $\widetilde{S}_i = S_i(v, \widetilde{B}_i)$ and let

$$Z = \{(\widetilde{w}, z) \in \widetilde{S}_i^{(0)} \times X^{(0)} | z \in X_w \setminus f_i((\widetilde{B}_i)_{\widetilde{w}})\}.$$

We define a relation $\overset{e}{\sim}$ on Z as follows:

$$(\widetilde{w}, z) \overset{e}{\sim} (\widetilde{w}', z') \text{ iff } (z = z' \text{ and } \langle \widetilde{w}, \widetilde{w}' \rangle \in \widetilde{B}_i^{(1)}) \ .$$

In order to define $\widetilde{B}_{i+1}$, we shall use the transitive closure $\overset{\overline{e}}{\sim}$ of the relation $\overset{e}{\sim}$. The rest of the proof relies on the following lemma.

Lemma 4.2. *If* $(\widetilde{w}_1, z) \overset{e}{\sim} (\widetilde{w}_2, z) \overset{e}{\sim} (\widetilde{w}_3, z) \overset{e}{\sim} (\widetilde{w}_4, z)$ *then there is* $(\widetilde{x}, z) \in Z$ *such that* $(\widetilde{w}_1, z) \overset{e}{\sim} (\widetilde{x}, z) \overset{e}{\sim} (\widetilde{w}_4, z)$.

Proof. By (P_i) and (Q_i), in $\widetilde{B}_{i-1}$ there are vertices $\widetilde{u}_j$ such that $\langle \widetilde{w}_j, \widetilde{u}_j, \widetilde{w}_{j+1} \rangle \in X$, $1 \leq j \leq 3$ and there are vertices $\widetilde{u'}_j$ such that $\widetilde{u'}_j \sim \widetilde{u}_j, \widetilde{w}_{j+1}, \widetilde{u}_{j+1}, j \in \{1, 2\}$ (possibly with $\widetilde{u'}_j = \widetilde{u}_{j+1}, j \in \{1, 2\}$). Let $u_j = f_i(\widetilde{u}_j), 1 \leq j \leq 3$ and let $u'_j = f_i(\widetilde{u'}_j), 1 \leq j \leq 2$ be vertices in X. By the (R_i) condition, we have $u'_j \sim u_j, w_{j+1}, u_{j+1}$ (possibly with $u'_j = u_{j+1}, j \in \{1, 2\}$). In order to apply 8-location, we have to analyze whether the homotopically trivial loop $(z, w_4, u_3, u'_2, u_2, u'_1, u_1, w_1)$ in X of length at most 8 is full.

In case $w_1 \sim w_3$, since $\widetilde{w}_1 \sim \widetilde{w}_2 \sim \widetilde{w}_3$, by the (R_i) condition, we have $\widetilde{w}_1 \sim \widetilde{w}_3$. Similarly, in case $w_2 \sim w_4$, by the (R_i) condition, we have $\widetilde{w}_2 \sim \widetilde{w}_4$. The lemma holds in these two cases trivially.

If $w_1 = w_4$, because $w_2 \sim w_3$, $w_3 \sim w_4$, and $w_2 \sim w_1$, we have $w_1, w_3 \in X_{w_2}$. Thus $w_1 \sim w_3$. Since $\widetilde{w}_1 \sim \widetilde{w}_2 \sim \widetilde{w}_3$, the (R_i) condition implies $\widetilde{w_1} \sim \widetilde{w_3}$. The lemma holds in this case trivially.



$w_1 = w_4$



$w_1 \sim w_4$, $\widetilde{u}_1 \sim \widetilde{u}_3$

Consider the case $w_1 \sim w_4$. To apply 8-location, we have to check whether the homotopically trivial loop of length at most 7 and containing at least the vertices w_4, u_3, u_1, w_1 is full. We have the cases $\widetilde{u}_1 = \widetilde{u}_3$; $\widetilde{u}_1 \sim \widetilde{u}_3$; $\widetilde{u}_1 = \widetilde{u}_2, \widetilde{u}_2 \sim \widetilde{u}_3$; $\widetilde{u}_1 \sim \widetilde{u}_2 \sim \widetilde{u}_3$; $\widetilde{u}_1 = \widetilde{u}_2, d(\widetilde{u}_2, \widetilde{u}_3) = 2$; $\widetilde{u}_1 \sim \widetilde{u}_2, d(\widetilde{u}_2, \widetilde{u}_3) = 2$; $d(\widetilde{u}_1, \widetilde{u}_2) = d(\widetilde{u}_2, \widetilde{u}_3) = 2$. We analyze the case $\widetilde{u}_1 \sim \widetilde{u}_3$. Except for the case $\widetilde{u}_1 = \widetilde{u}_3$ (which will be treated separately in the next paragraph), the other cases can be treated similarly. Assume $w_1 \nsim u_3, w_4 \nsim u_1$ (if $w_1 \sim u_3$ or $w_4 \sim u_1$, we have reached case $\widetilde{u}_1 = \widetilde{u}_3$ treated below). Then the homotopically trivial loop $\gamma = (w_4, u_3, u_1, w_1)$ is full. By 8-location there is a vertex $x \in X$ adjacent to all vertices of γ. So $\langle x, u_1 \rangle$ and $\langle x, u_3 \rangle$ are edges in X. By the (R_i) condition applied to the vertices $\widetilde{u}_1$ and $\widetilde{u}_3$, there is a vertex $\widetilde{x}$ in $\widetilde{B}_i$ such that $f_i(\widetilde{x}) = x$ while $\langle \widetilde{x}, \widetilde{u}_1 \rangle$ and $\langle \widetilde{x}, \widetilde{u}_3 \rangle$ are edges in $\widetilde{B}_i$. By the (R_i) condition, since $x \sim w_4$, we have $\widetilde{x} \sim \widetilde{w}_4$. Similarly we get $\widetilde{x} \sim \widetilde{w}_1$. Finally, because $w_1 \sim w_4$ and $\widetilde{w}_1$ and $\widetilde{w}_4$ both belong to the link of $\widetilde{x}$ in $\widetilde{B}_i$, the (R_i) condition implies $\widetilde{w}_1 \sim \widetilde{w}_4$. The lemma holds in this case trivially.

Consider further the case when $w_1 \sim w_4$ and $\widetilde{u}_1 = \widetilde{u}_3$. Since $\widetilde{w}_1, \widetilde{w}_4 \in (\widetilde{B}_i)_{\widetilde{u}_1}$ and $w_1 \sim w_4$, the (R_i) condition implies $\widetilde{w}_1 \sim \widetilde{w}_4$. The lemma holds in this case trivially.



$w_1 \sim w_4$; $\widetilde{u}_1 = \widetilde{u}_3$

From now on assume $w_1 \nsim w_3$, $w_2 \nsim w_4$, $w_1 \neq w_4$ and $w_1 \nsim w_4$.

Suppose $z = u_1$. By the definition of the set Z, $z \in X_{w_1}$. Since $z = u_1$ and $u_1 \sim w_1$, by the (R_i) condition applied to the vertex $\widetilde{w}_1$, there is in $\widetilde{B}_i$ a vertex $\widetilde{z}$ such that $\widetilde{z} \sim \widetilde{w}_1$ and $f_i(\widetilde{z}) = z$. So $\langle \widetilde{z}, \widetilde{w}_1 \rangle \in \widetilde{B}_i$. But since $(\widetilde{w}_1, z) \in Z$, $\langle \widetilde{z}, \widetilde{w}_1 \rangle \notin \widetilde{B}_i$. Since we have reached a contradiction, $z \neq u_1$.

Suppose $z \sim u_1$. By the definition of the set Z, $z \in X_{w_1}$. By the (R_i) condition applied to the vertex $\widetilde{u}_1$, there is in $\widetilde{B}_i$ a vertex $\widetilde{z}$ such that $\widetilde{z} \sim \widetilde{u}_1$ and $f_i(\widetilde{z}) = z$. So because $\widetilde{w}_1$ and $\widetilde{z}$ are both in the link of $\widetilde{u}_1$ in $\widetilde{B}_i$, we have, by the (R_i) condition, $\langle \widetilde{z}, \widetilde{w}_1 \rangle \in \widetilde{B}_i$. But since $(\widetilde{w}_1, z) \in Z$, $\langle \widetilde{z}, \widetilde{w}_1 \rangle \notin \widetilde{B}_i$. This yields a contradiction and hence $z \nsim u_1$.

In conclusion the homotopically trivial loop $(z, w_4, u_3, u_2', u_2, u_1', u_1, w_1)$ is full. Because its length is at most 8, we may apply 8-location. There are 7 cases to be analyzed: $\widetilde{u}_1 = \widetilde{u}_3$; $\widetilde{u}_1 \sim \widetilde{u}_2$, $\widetilde{u}_2 = \widetilde{u}_3$; $\widetilde{u}_1 \sim \widetilde{u}_3$; $\widetilde{u}_1 \sim \widetilde{u}_2 \sim \widetilde{u}_3$; $d(\widetilde{u}_1, \widetilde{u}_2) = 2, \widetilde{u}_2 = \widetilde{u}_3$; $\widetilde{u}_1 \sim \widetilde{u}_2, d(\widetilde{u}_2, \widetilde{u}_3) = 2$; $d(\widetilde{u}_1, \widetilde{u}_2) = d(\widetilde{u}_2, \widetilde{u}_3) = 2$.

Case 1 is when $\widetilde{u}_1 = \widetilde{u}_3$. We obtain in X, by the (R_i) condition and the definition of the set Z, a full homotopically trivial loop $\gamma = (z, w_4, u_1, w_1)$ of length 4. By 8-location, the loop γ is contained in the link of a vertex x. Hence, by the (R_i) condition applied to the vertex $\widetilde{u_1}$, there exists in $\widetilde{B_i}$ a vertex $\widetilde{x}$ such that $\widetilde{x} \sim \widetilde{u_1}$ and $f_i(\widetilde{x}) = x$. Moreover, according to the (R_i) condition, the vertices $\widetilde{w_4}$ and $\widetilde{w_1}$ are adjacent to $\widetilde{x}$ in $\widetilde{B_i}$. Hence, once we show that $(\widetilde{x}, z) \in Z$, the lemma is in this case proven. Assume that $(\widetilde{x}, z) \notin Z$. Then by the (R_i) condition, there exists a vertex $\widetilde{z} \in \widetilde{B_i}$ such that $\langle \widetilde{z}, \widetilde{x} \rangle \in \widetilde{B_i}$. Because $\widetilde{w_1}$ and $\widetilde{z}$ both belong to the link of $\widetilde{x}$ in $\widetilde{B_i}$, this implies that $\langle \widetilde{z}, \widetilde{w_1} \rangle \in \widetilde{B_i}$. But, since $(\widetilde{w_1}, z) \in Z$, we have $\langle \widetilde{z}, \widetilde{w_1} \rangle \notin \widetilde{B_i}$. So, because we have reached a contradiction, $(\widetilde{x}, z) \in Z$.



Case 1

Case 2 is when $\widetilde{u}_1 \sim \widetilde{u}_2$, $\widetilde{u}_2 = \widetilde{u}_3$. We obtain in X, by the (R_i) condition and the definition of the set Z, a homotopically trivial loop $\gamma = (z, w_4, u_2, u_1, w_1)$ of length 5. If $\widetilde{w}_1 \sim \widetilde{u}_2$ or $\widetilde{w}_4 \sim \widetilde{u}_1$, we are in case 1 treated above. If this does not happen, the loop γ is full. Because X is 8-located, γ is contained in the link of a vertex x. Hence, by the (R_i) condition applied to the vertex $\widetilde{u_2}$, there exists in $\widetilde{B_i}$ a vertex $\widetilde{x}$ such that $\widetilde{x} \sim \widetilde{u_2}$ and $f_i(\widetilde{x}) = x$. Moreover, according to the (R_i) condition, the vertices $\widetilde{w_4}$ and $\widetilde{w_1}$ are adjacent to $\widetilde{x}$ in $\widetilde{B_i}$. Hence, if $(\widetilde{x}, z) \in Z$, the lemma is in this case proven. We show that $(\widetilde{x}, z) \in Z$ as in case 1. Note that the case $\widetilde{u}_1 \sim \widetilde{u}_3$ can be treated similarly.



Case 2



$\widetilde{u}_1 \sim \widetilde{u}_3$

Case 3 is when $\widetilde{u}_1 \sim \widetilde{u}_2 \sim \widetilde{u}_3$. We obtain in X, by the (R_i) condition and the definition of the set Z, a homotopically trivial loop $\gamma = (z, w_4, u_3, u_2, u_1, w_1)$ of length 6. If $\widetilde{w}_1 \sim \widetilde{u}_3$, or $\widetilde{w}_1 \sim \widetilde{u}_2$ and $\widetilde{w}_4 \sim \widetilde{u}_2$, or $\widetilde{w}_4 \sim \widetilde{u}_1$, we have reached case 1 treated above. If $\widetilde{w}_1 \sim \widetilde{u}_2$ and $\widetilde{w}_4 \nsim \widetilde{u}_2$, or $\widetilde{w}_4 \sim \widetilde{u}_2$ and $\widetilde{w}_1 \nsim \widetilde{u}_2$, we are in case 2 treated above. If none of these situations occur, the loop is full. Because X is 8-located, γ is contained in the link of a vertex x. Hence, by the (R_i) condition applied to the vertex $\widetilde{u_2}$, there exists in $\widetilde{B_i}$ a vertex $\widetilde{x}$ such that $\widetilde{x} \sim \widetilde{u_2}$ and $f_i(\widetilde{x}) = x$. Moreover, according to the (R_i) condition, the vertices $\widetilde{w_4}$ and $\widetilde{w_1}$ are adjacent to $\widetilde{x}$ in $\widetilde{B_i}$. Hence, because $(\widetilde{x}, z) \in Z$, the lemma is in this case proven. We show that $(\widetilde{x}, z) \in Z$ as in case 1. Note that the case $d(\widetilde{u}_1, \widetilde{u}_2) = 2, \widetilde{u}_2 = \widetilde{u}_3$ can be treated similarly.



Case 3



$d(\widetilde{u}_1, \widetilde{u}_2) = 2, \widetilde{u}_2 = \widetilde{u}_3$

Case 4 is when $\widetilde{u}_1 \sim \widetilde{u}_2, d(\widetilde{u}_2, \widetilde{u}_3) = 2$. We obtain in X, by the (R_i) condition and the definition of the set Z, a homotopically trivial loop $\gamma = (z, w_4, u_3, u'_2, u_2, u_1, w_1)$ of length 7. If $\widetilde{w}_1 \sim \widetilde{u}_3$, or $\widetilde{w}_4 \sim \widetilde{u}_1$, or $\widetilde{w}_1 \sim \widetilde{u}'_2$ and $\widetilde{w}_4 \sim \widetilde{u}'_2$, or $\widetilde{w}_1 \sim \widetilde{u}_2$ and $\widetilde{w}_4 \sim \widetilde{u}_2$, we are in case 1 treated above. If $\widetilde{w}_1 \sim \widetilde{u}'_2$ but $\widetilde{w}_4 \nsim \widetilde{u}'_2$, or $\widetilde{w}_1 \sim \widetilde{u}_2$ and $\widetilde{w}_4 \sim \widetilde{u}'_2$, we have reached case 2 treated above. If $\widetilde{w}_1 \sim \widetilde{u}_2$ but $\widetilde{w}_4 \nsim \widetilde{u}'_2$ and $\widetilde{w}_4 \nsim \widetilde{u}_2$, we have reached case 3 treated above. If none of these situations occur, the loop γ is full. Because X is $8-$located, γ is contained in the link of a vertex x. Hence, by the (R_i) condition applied to the vertex $\widetilde{u_2}$, there exists in $\widetilde{B_i}$ a vertex $\widetilde{x}$ such that $\widetilde{x} \sim \widetilde{u_2}$ and $f_i(\widetilde{x}) = x$. Moreover, according to the (R_i) condition, the vertices $\widetilde{w_4}$ and $\widetilde{w_1}$ are adjacent to $\widetilde{x}$ in $\widetilde{B_i}$. Hence, because $(\widetilde{x}, z) \in Z$, the lemma is in this case proven.

Case 5 is when $d(\widetilde{u}_1, \widetilde{u}_2) = d(\widetilde{u}_2, \widetilde{u}_3) = 2$. We obtain in X, by the (R_i) condition and the definition of the set Z, a homotopically trivial loop $\gamma = (z, w_4, u_3, u'_2, u_2, u'_1,$



Case 4

$u_1, w_1)$ of length 8. If $\widetilde{w_1} \sim \widetilde{u}_1'$ and $\widetilde{w_4} \sim \widetilde{u}_1'$, we are in case 1 treated above. If $\widetilde{w_4} \sim \widetilde{u}_1'$ and $\widetilde{w_1} \nsim \widetilde{u}_1'$, we have reached case 2 treated above. If $\widetilde{w_4} \sim \widetilde{u}_2$ but $\widetilde{w_1} \nsim \widetilde{u}_2$, $\widetilde{w_1} \nsim \widetilde{u}_1'$ and $\widetilde{w_4} \nsim \widetilde{u}_1'$, we have reached case 3 treated above. If $\widetilde{w_1} \sim \widetilde{u}_1'$, but $\widetilde{w_1} \nsim \widetilde{u_2}$, $\widetilde{w_1} \nsim \widetilde{u}_2'$, $\widetilde{w_1} \nsim \widetilde{u}_3$, $\widetilde{w_4} \nsim \widetilde{u}_2'$, $\widetilde{w_4} \nsim \widetilde{u_2}$, $\widetilde{w_4} \nsim \widetilde{u}_1'$, we have reached case 4 treated above. If none of these situations occur, the loop γ is full. Because X is 8-located, γ is contained in the link of a vertex x. Hence, by the (R_i) condition applied to the vertex $\widetilde{u_2}$, there exists in $\widetilde{B_i}$ a vertex $\widetilde{x}$ such that $\widetilde{x} \sim \widetilde{u_2}$ and $f_i(\widetilde{x}) = x$. Moreover, according to the (R_i) condition, the vertices $\widetilde{w_4}$ and $\widetilde{w_1}$ are adjacent to $\widetilde{x}$ in $\widetilde{B_i}$. Hence, because $(\widetilde{x}, z) \in Z$, the lemma is in this case proven.



Case 5

□

According to the previous lemma, if $(\widetilde{u}, z) \overset{\overline{e}}{\sim} (\widetilde{w}, z)$, then there is a vertex $\widetilde{x} \in \widetilde{S}_i$ such that $(\widetilde{x}, z) \in Z$ and $\langle \widetilde{x}, \widetilde{u} \rangle, \langle \widetilde{x}, \widetilde{w} \rangle \in \widetilde{B}_i$.

We define further the flag simplicial complex $\widetilde{B}_{i+1}$. Its 0-skeleton is defined as the set $\widetilde{B}_{i+1}^{(0)} = \widetilde{B}_i^{(0)} \cup (Z/ \overset{\overline{e}}{\sim})$. Further we define the 1-skeleton $\widetilde{B}_{i+1}^{(1)}$ of $\widetilde{B}_{i+1}$. Edges between vertices of $\widetilde{B}_i$ are the same as in $\widetilde{B}_i$. For every $\widetilde{w} \in \widetilde{S}_i^{(0)}$, there are edges joining $\widetilde{w}$ with $[\widetilde{w}, z] \in Z/ \overset{\overline{e}}{\sim}$ (here $[\widetilde{w}, z]$ denotes the equivalence class of $(\widetilde{w}, z) \in Z$) and there are edges joining $\langle \widetilde{w}, z \rangle$ with $\langle \widetilde{w}, z' \rangle$, for $\langle z, z' \rangle \in X$. Once we have defined the 1-skeleton of $\widetilde{B}_{i+1}$, the higher dimensional skeleta are determined by the flagness property.

The map $f_{i+1} : \widetilde{B}_{i+1}^{(0)} \to X$ is defined by $f_{i+1}|_{\widetilde{B}_i} = f_i$ and $f_{i+1}(\langle \widetilde{w}, z \rangle) = z$. As proven in [Osa15a] (Theorem 3.4), this map can be extended simplicially. By the simplicial extension, we can define the map $f_{i+1} : \widetilde{B}_{i+1} \to X$. The proof of the conditions (P_{i+1}), (Q_{i+1}) and (R_{i+1}) uses the above lemma and it is similar to the one given in [Osa15a] (Theorem 3.4).

So we have built inductively a complex $\widetilde{X} = \cup_{i=1}^{\infty} \widetilde{B}_i$ which satisfies the $SD'_n(O)$ property for each n. Inductively we have also constructed a map $f = \cup_{i=1}^{\infty} f_i : \widetilde{X} \to X$ which is a covering map. Because $\widetilde{X}$ was built such that it satisfies, for each n, the $SD'_n(O)$ property, it is, by Proposition 2.1, simply connected. So $\widetilde{X}$ is the universal cover of X. Because the universal cover of X is unique and since the vertex O is arbitrary, $\widetilde{X}$ satisfies the $SD'_n(O)$ property for each vertex O and for each natural number n. Hence we have constructed the universal cover of X which satisfies the SD' property.

□

Theorem 3.2 and Theorem 4.1 imply the paper's main result.

Theorem 4.3. *Let X be a simply connected, 8-located simplicial complex. Then the $0-$skeleton of X with a path metric induced from $X^{(1)}$, is $\delta-$hyperbolic, for a universal constant δ.*

References

[BCC+13] B. Brešar, J. Chalopin, V. Chepoi, T. Gologranc, and D. Osajda, *Bucolic complexes*, Adv. Math. **243** (2013), 127–167.

[CCHO14] J. Chalopin, V. Chepoi, H. Hirai, and D. Osajda, *Weakly modular graphs and nonpositive curvature* (2014), preprint, available at `arXiv:1409.3892`.

[CO15] V. Chepoi and D. Osajda, *Dismantlability of weakly systolic complexes and applications*, Trans. Amer. Math. Soc. **367** (2015), no. 2, 1247–1272.

[GO14] R. Gómez-Ortells, *Compactly supported cohomology of systolic 3-pseudomanifolds*, Colloq. Math. **135** (2014), no. 1, 103–112.

[Gro87] M. Gromov, *Hyperbolic groups*, Essays in group theory, Math. Sci. Res. Inst. Publ., vol. 8, Springer, New York, 1987, pp. 75–263.

[Hag03] F. Haglund, *Complexes simpliciaux hyperboliques de grande dimension*, Prepublication Orsay **71** (2003), preprint, available at `http://www.math.u-psud.fr/~haglund/cpl_hyp_gde_dim.pdf`.

[JŚ03] T. Januszkiewicz and J. Świątkowski, *Hyperbolic Coxeter groups of large dimension*, Comment. Math. Helv. **78** (2003), no. 3, 555–583.

[JŚ06] ———, *Simplicial nonpositive curvature*, Publ. Math. Inst. Hautes Études Sci. **104** (2006), 1–85.

[JŚ07] ———, *Filling invariants of systolic complexes and groups*, Geom. Topol. **11** (2007), 727–758.

[Osa07] D. Osajda, *Connectedness at infinity of systolic complexes and groups*, Groups Geom. Dyn. **1** (2007), no. 2, 183–203.

[Osa08] ———, *Ideal boundary of 7-systolic complexes and groups*, Algebr. Geom. Topol. **8** (2008), no. 1, 81–99.

[Osa13a] ———, *A construction of hyperbolic Coxeter groups*, Comment. Math. Helv. **88** (2013), no. 2, 353–367.

[Osa13b] ———, *A combinatorial non-positive curvature I: weak systolicity* (2013), preprint, available at `arXiv:1305.4661`.

[Osa15a] ———, *Combinatorial negative curvature and triangulations of 3-manifolds*, Indiana Univ. Math. J. **64** (2015), no. 3, 943–956.

[Osa15b] ———, *Normal subgroups of SimpHAtic groups* (2015), submitted, available at `arXiv:1501.00951`.

[OŚ15] D. Osajda and J. Świątkowski, *On asymptotically hereditarily aspherical groups*, Proc. London Math. Soc., posted on 2015, DOI 10.1112/plms/pdv021, (to appear in print).

[Pap95] P. Papasoglu, *Strongly geodesically automatic groups are hyperbolic*, Invent. Math. **121** (1995), no. 2, 323–334, DOI 10.1007/BF01884301. MR1346209 (96h:20073)